PEPTIDEDOSAGE

Thyreogen Dosage Chart

Medically reviewed June 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

1-2 mg/day SC (educational); 10 mg/capsule oral

20 mg vial + 2 mL bacteriostatic water

10 mg/mL · 1 unit = 100 mcg on a U-100 syringe

PHASE	DOSE PER INJECTION	UNITS (PER INJECTION)
Microdose intro (days 1-3)	500 mcg	5 units (0.05 mL)
Conservative course (days 1-10)	1000 mcg (1 mg)	10 units (0.10 mL)
Standard course (days 1-10)	2000 mcg (2 mg)	20 units (0.20 mL)

Thyreogen dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	20 MG VIAL + 2 ML (10 MG/ML)
500 mcg	5 units (0.05 mL)
1 mg	10 units (0.1 mL)
2 mg	20 units (0.2 mL)

STORAGE

Capsules: store in a cool, dry place at about 15-25 °C, away from light and moisture. For the educational lyophilized-vial model: store at -20 °C long term, or 2-8 °C for short periods, protected from light; once reconstituted, keep refrigerated at 2-8 °C and use within about 3-4 weeks.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.