PEPTIDEDOSAGE

Vilon Dosage Chart

Medically reviewed May 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

67–670 mcg once daily for 5 consecutive days per cycle

20 mg vial + 3 mL bacteriostatic water

6.67 mg/mL · 1 unit = 66.7 mcg on a U-100 syringe

PHASE / CYCLE	DAILY DOSE (MCG)	UNITS (PER INJECTION) (ML)
Cycle 1, Day 1	67 mcg (0.067 mg)	1 unit (0.01 mL)
Cycle 1, Day 2	133 mcg (0.133 mg)	2 units (0.02 mL)
Cycle 1, Day 3	200 mcg (0.20 mg)	3 units (0.03 mL)
Cycle 1, Day 4	267 mcg (0.267 mg)	4 units (0.04 mL)
Cycle 1, Day 5	333 mcg (0.33 mg)	5 units (0.05 mL)
Cycle 2+ (Days 1–5)	333–667 mcg (0.33–0.67 mg)	5–10 units (0.05–0.10 mL)

Vilon dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	20 MG VIAL + 3 ML (6.67 MG/ML)
67 mcg	1 units (0.01 mL)
133 mcg	2 units (0.02 mL)
200 mcg	3 units (0.03 mL)
267 mcg	4 units (0.04 mL)
333 mcg	5 units (0.05 mL)

STORAGE

Lyophilized frozen; reconstituted refrigerated; avoid repeated freeze–thaw.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.