PEPTIDEDOSAGE

Visoluten Dosage Chart

Medically reviewed June 21, 2026 · Page updated November 21, 2026

TYPICAL RESEARCH DOSE

2.5-5 mg/day SC reference; 10-20 mg/day oral (2 caps)

20 mg vial + 2 mL bacteriostatic water

10 mg/mL · 1 unit = 100 mcg on a U-100 syringe

PHASE	DOSE PER INJECTION	UNITS (PER INJECTION)
Conservative course (days 1-10)	2500 mcg (2.5 mg)	25 units (0.25 mL)
Standard course (days 1-10)	5000 mcg (5 mg)	50 units (0.50 mL)

Visoluten dose to syringe units, by vial size

U-100 insulin syringe: 100 units = 1 mL.

DOSE	20 MG VIAL + 2 ML (10 MG/ML)
2.5 mg	25 units (0.25 mL)
5 mg	50 units (0.5 mL)

STORAGE

Lyophilized/capsule material: store at -20 °C long term, or 2-8 °C for short periods, protected from light and moisture. Once reconstituted for the educational reference, keep refrigerated at 2-8 °C and use within about 3-4 weeks; capsules should be kept dry at room temperature per the supplier label.

Educational reference only, not medical advice. Figures restate the research protocol published at the address above.